



臺北醫學大學 泌尿腎臟研究中心 會議記錄

時間：**112年4月21日(星期五) 14:00-15:00**

地點：視訊會議-(請以正式全名登入會議室，以利進行會議簽到)

使用 Google Meet (會議前 10 分鐘即開啟會議室)

會議室連結：<https://meet.google.com/xmk-tzqg-dca>

(敬略稱位)

會議主席：吳麥斯

與會人員：

【附醫】劉明哲、葉劭德、吳建志、林孝友、吳政誠、張景欣、陳偉傑、顧芳瑜、
羅詩修、戴定恩、方德昭、陳錫賢、林彥仲、吳岳霖、高治圻、陳靜怡、
葉曙慶、邵月珠

【萬芳】溫玉清、李良明、林克勳、林雍偉、蕭志豪、許軒豪、賴宗豪、鍾卓興、
鄭仲益、陳作孝、蘇裕謀、劉崇德、楊韻紅、李明哲、鍾卓興

【雙和】吳佳璋、陳冠州、劉家宏、江怡德、鄒凱亦、高偉棠、胡書維、魏汶玲、
吳美儀、洪麗玉、鄭彩梅、邱怡仁、陳佑瑋、廖家德、游博翰、陳正憲、
邱惠雯


【新國民】許永和、鄒居霖

長官指導：

林建煌校長、李岡遠研發長、許志成教授、崔克宏副院長、陳瑞明所長

議程：

一、慢性腎病團隊、整合透析介入照護團隊 小組報告



慢性腎病團隊

萬芳: 鄭仲益、蘇裕謀

北醫: 林彥仲、葉曜慶

雙和: 鄭彩梅、廖家德、宋立勤

新國民: 許永和、鄧居霖

數位醫療、遠距監控、個人精準、腎臟再生

報告人：鄭仲益醫師

112.04.21

上期研究追蹤



- Artificial intelligence in predicting kidney interstitial fibrosis and tubular atrophy severity
- Status: Manuscript in submission
Corresponding author: Yen-Chung Lin

3



Clinical and pathological features
of ANCA associated GN before
and after COVID-19 epidemic

5

ANCA glomerulonephritis & vasculitis (ANCA-GN & AAV)



- ANCA: Anti-Neutrophil Cytoplasmic Antibody directly against cytoplasmic antigen.
- Necrotizing vasculitis affecting small vessels
- Myeloperoxidase (MPO-ANCA) mostly **p-ANCA**
- Proteinase 3 (PR3-ANCA) mostly **c-ANCA**

ANCA renal-limited vasculitis (RLV)

ANCA-associated vasculitis (AAV)

1. Microscopic polyangiitis (MPA)

2. Granulomatosis with polyangiitis (Wegener)

3. Eosinophilic granulomatosis with polyangiitis (Churg-Strauss)


CHCC 2012 categories of ANCA-associated vasculitis 6

ANCA-GN



- 50% risk of death or renal failure at 5 yrs when eGFR < 50 ml/min.
- Often related to systemic vasculitis (AAV), MPA, GPA or RLV
- **RPGN**, subnephrotic range proteinuria, microhematuria, and hypertension over days to months.
- **Variable clinical courses:**
 1. Asymptomatic urinary abnormalities
 2. Indolent, remitting, and relapsing course
 3. Chronic remitting course, but cause chronic progressive renal failure
 4. Slowly progressive course for several years.

Pathogenic model highlighting the differences between PR3-ANCA and MPO-ANCA vasculitis.



ANCA-associated glomerulonephritis (ANCA-GN)



- 20/million population
- Most common form of new-onset GN in adult > 50 years, peak 60- to 70- years
- More common in white and Asian
- Different spectrum of clinicopathologic phenotypes and serotypes of ANCA disease.
- Reported of HLA-DR4 and DRB4 as potential risk alleles for SARS-CoV2 vaccine-related ANCA-associated GN

Table 1. ANCA associated GN in Taiwan (2001-2016)^a

TP series: bimonthly (15 years)	3.0%*
2001-2007	2.1% (7/326)
2008-2016.09	3.8% (13/340)
TMU series: monthly (4 years)	3.5%*
2012-2016.10	3.5% (8/229)
TC series: bimonthly (15 years)	11.7%*
2001-2007	8.9% (11/123)
2008-16.10	13.4% (27/202)

* $P < 0.0001$ (TP v. TC); Geographic variation? Environment?
^a Six anti-GBM/P-ANCA dual (+) cases are not included.

許輝吉 教授 整理資料

Proposal to categorize the glomerular lesions linked to treatment response:

Unrescuable:

Acute: Severe glomerular necrosis (NGN), global or circumferential to massive crescents (GC)

Chronic: Global sclerosis (GS)

Rescuable: Potential recovery from the re-expansion or preserved tufts after the halt of disease activity. (proven by rebiopsy)

Acute : FN and FC

Chronic: FS

許輝吉 教授 整理資料

	ANCA	P-ANCA	C-ANCA	Atypical P-ANCA
Case no.	88	61 (1 double positive, anti-GBM and P-ANCA)	20	7
Renal biopsied	13	11	2	0
Female (%)	42(47.7%)	33 (53.2%)	6 (30%)	3 (42.9%)
Age (yrs)	66.5±15.5	67.0±15.6	62.9±15.2	72.0±15.7
Bodyweight (kg)	60.0±16.3	59.0±14.0	61.7±20.4	64.4±23.5
BMI	23.5±5.8	23.1±4.4	23.7±7.4	26.0±10.3
BUN (mg/dL)	36.6±33.4	34.3±28.3	48.8±47.1	22.1±18.1
Cr (mg/dL)	2.3±2.6	2.5±2.8	2.3±2.1	1.0±0.8

14

	ANCA-GN	Before COVID-19	After COVID-19
Renal biopsied	13	7	6
Female (%)	8 (61.5%)	4 (57.1%)	4 (66.7%)
Age (yrs)	63.9±10.9	64.6±13.1	63.2±8.9
Bodyweight (kg)	56.7±8.6	56.4±7.2	57.0±10.8
BMI	21.7±2.9	20.8±2.4	22.7±3.4
BUN	66.1±27.3	69.9±26.4	61.7±30.2
Cr	5.0±3.3	5.1±3.7	4.8±3.2
Manchester ANCA Renal Risk Score	7.2±2.3	7.3±2.7	7.2±1.9

Normal glomeruli < 10% 10-25% > 25%

Tubular atrophy/interstitial fibrosis ≤ 25% > 25%


Renal function eGFR (ml/min): > 15 ≤ 15



<https://bunes.apple.com/us/app/anca-score/id1415367084?i=de&ls=1&mt=8>

15

Percentage of patients without dialysis or death



16

Future work



- Recruit TMUH and SHH data
- IgA and MGN before and after COVID-19 era
- Genetic predisposition, HLA DR4 and HLA DRB4

17




Progress Report

雙和醫院 腎臟內科 邱怡仁
整合介入性照護團隊
APR-21, 2023

Blood flow plays a role in AVG patency

- **Thrombosis** was one the main reason for the graft patency loss, which leads to increasing medical expense for rescue of the vascular access
- KDOQI (Kidney Disease Outcomes Quality Initiative) guidelines for vascular access suggest AVG surveillance by **measuring access blood flow** is supplementary to regular clinical monitoring, to **improve AVG patency**

Decreased blood flow in prosthetic vessels is associated with thrombosis



- Intimal hyperplasia over the venous-graft junction
- Systemic hemodynamics
Blood pressure
Cardiac output


J Vasc Access. 2020 Mar;21(2):195-203.



- KDOQI guidelines suggested arm exercise may be beneficial to vascular access maturation
- Scarce evidence focusing on **the impact of the arm posture** on the blood flow of the AVG



Method



49 HD patients using *forearm AVG* > 4 weeks from SHH dialysiscenter (IRB-N202104038; 2022/04-2023/01)

Four different position

A Sitting extension (rest) B Sitting flexion
C Lying extension (rest) D Lying flexion


Philips EPIQ 7 diagnostic ultrasound (USA) and an L12-3MHz Ultrasound transducer (WA, USA) and 60-degree Doppler angle, also same graft material

Preliminary results




Variables	all patients (n = 49)
Age (years)	65.96 ± 11.28
Sex	
Male (%)	22 (44.9)
female (%)	27 (55.1)
BMI	25.93 ± 6.87
Diabetes mellitus (%)	22 (44.9)
Hypertension (%)	36 (73.5)
Current smoker (%)	5 (10.2)
Coronary artery disease (%)	15 (30.6)
Cerebrovascular accident (%)	4 (8.2)
Peripheral vascular disease (%)	5 (10.2)
Graft sided (%)	
left	43 (87.8)
right	6 (12.2)
Graft configuration (%)	
loop	45 (91.8)
straight	4 (8.2)
Cross elbow (%)	
crossed	12 (24.5)
not crossed	37 (75.5)
Graft age (months), median (IQR)	15.00 (47.00)
History of graft thrombosis (%)	24 (49.0)
Systolic BP (mmHg)	149.59 ± 25.65
Diastolic BP (mmHg)	69.92 ± 17.94
Heart rate (beats per min)	79.45 ± 15.05


Sitting position does **not alter** AVG blood flow



Lying with arm flexion **reduces** AVG blood flow



Systolic BP may **not** correlate with AVG blood flow



Future works



- Subgroup analysis
 - T2DM
 - Age (65Y)
 - Intervention